



GPDs with CLAS12

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

F.-X. Girod
April 12th 2011

1 Introduction

2 CLAS 12

3 Proposed measurements

Proton DVCS

Neutron DVCS

Other highlights

4 GPD extraction

5 Summary

Introduction

Momentum distributions in the transverse plane

$$q_X(x, \vec{b}_\perp) = \int \frac{d^2 \vec{\Delta}_\perp}{(2\pi)^2} H(x, 0, t) e^{-i \vec{\Delta}_\perp \cdot \vec{b}_\perp} - \frac{1}{2M} \frac{\partial}{\partial b_y} \int \frac{d^2 \vec{\Delta}_\perp}{(2\pi)^2} E(x, 0, t) e^{-i \vec{\Delta}_\perp \cdot \vec{b}_\perp}$$

M. Burkardt, Phys. Rev. **D62**, (2000) 071503

$\xi \neq 0$ in M. Diehl, Eur. Phys. J. **C25** (2002) 223

Energy-momentum tensor of q flavored quarks

$$\langle p_2 | \hat{T}_{\mu\nu}^q | p_1 \rangle = \tilde{U}(p_2) \left[M_2^q(t) \frac{P_\mu P_\nu}{M} + J^q(t) \frac{i(P_\mu \sigma_{\nu\rho} + P_\nu \sigma_{\mu\rho}) \Delta^\rho}{2M} + d_1^q(t) \frac{\Delta_\mu \Delta_\nu - g_{\mu\nu} \Delta^2}{5M} \right] U(p_1)$$

To measure gravitational FFs : **graviton** scattering or **GPDs** identities :

$$J^q(t) = \frac{1}{2} \int_{-1}^1 dx x [H^q(x, \xi, t) + E^q(x, \xi, t)] \quad , \quad M_2^q(t) + \frac{4}{5} d_1(t) \xi^2 = \frac{1}{2} \int_{-1}^1 dx x H^q(x, \xi, t)$$

(Ji's sum rule)

Physical content of GPDs : Energy-momentum tensor of q flavored quarks


$$\langle p_2 | \hat{T}_{\mu\nu}^q | p_1 \rangle = \bar{U}(p_2) \left[M_2^q(t) \frac{P_\mu P_\nu}{M} + J^q(t) \frac{i(P_\mu \sigma_{\nu\rho} + P_\nu \sigma_{\mu\rho}) \Delta^\rho}{2M} + d_1^q(t) \frac{\Delta_\mu \Delta_\nu - g_{\mu\nu} \Delta^2}{5M} \right] U(p_1)$$

$M_2(t) \longleftrightarrow T_{00}$: mass distributions inside the hadron

$J(t) \longleftrightarrow T_{0i}$: angular momentum distributions

$d_1(t) \longleftrightarrow T_{ij}$: forces and pressure distributions

Large N_c χ QSM from K.Goeke *et al.* in Phys. Rev. **D75** (2007) 094021



$$\text{Stability} \Rightarrow \int_0^\infty dr r^2 p(r) = 0$$

$r < 0.57 \text{ fm} \Rightarrow p(r) > 0 \leftrightarrow$ **repulsion** (quark core)

$r > 0.57 \text{ fm} \Rightarrow p(r) < 0 \leftrightarrow$ **attraction** (pion cloud)

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

Access to GPDs : the DVCS process

Observables in the Bjorken limit

$$\gamma^* p \rightarrow \gamma p'$$

Bjorken regime :


$$Q^2 \rightarrow \infty,$$

$$\nu \rightarrow \infty,$$

$$x_B = Q^2 / 2M\nu \text{ fixed}$$

$$\left(\xi \rightarrow \frac{x_B}{2-x_B} \right)$$

Müller, Ji, Radyushkin (1994-96)



$$ep \rightarrow ep\gamma$$

$$\sigma(ep \rightarrow ep\gamma) \propto \left| \text{DVCS} + \text{BH} \right|^2$$

(a) (b) (c)

Diehl, Gousset, Pire, Ralston (1997)

Belitsky, Müller, Kirchner (2002-10)



$$A_{LU} = \frac{d^4\sigma^{\rightarrow} - d^4\sigma^{\leftarrow}}{d^4\sigma^{\rightarrow} + d^4\sigma^{\leftarrow}} \stackrel{\text{twist-2}}{\approx} \frac{\alpha \sin \phi + \dots}{1 + \beta \cos \phi + \dots}$$

$$\alpha \propto \left(F_1 \mathcal{H} + \xi G_M \tilde{\mathcal{H}} - \frac{t}{4M^2} F_2 \mathcal{E} \right)$$

$$\mathcal{H}(\xi, t) = \pi \sum_q Q_q^2 [H^q(\xi, \xi, t) - H^q(-\xi, \xi, t)]$$

$$A_{UL} \propto \left(F_1 \tilde{\mathcal{H}} + \xi G_M \mathcal{H} + G_M \frac{\xi}{1+\xi} \mathcal{E} + \dots \right) \sin \phi$$

Observables sensitivities to GPD



	Im	Re
\mathcal{H}	A_{LU}	σ, A_{LL}
$\tilde{\mathcal{H}}$	A_{UL}	
\mathcal{E}	$A_{\text{UT}}, A_{\text{LT}}$	

DVCS

	Meson	Flavor
$\tilde{\mathcal{H}}, \tilde{\mathcal{E}}$	π^+	$\Delta u - \Delta d$
	π^0	$2\Delta u + \Delta d$
	η	$2\Delta u - \Delta d + 2\Delta s$
\mathcal{H}, \mathcal{E}	ρ^+	$u - d$
	ρ^0	$2u + d$
	ω	$2u - d$
	ϕ	s

DVMP

Only a global analysis of all observables can disentangle GPDs

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS


Neutron DVCS

Other highlights

GPD extraction

Summary

12 GeV upgrade



Upgraded apparatus

Generalized Parton
Distributions with
CLAS12

F.-X. Girod

Experimental Hall B

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS



Other highlights

GPD extraction

Summary






Higher energy, luminosity, hermiticity, analyzing power

HTCC




CTOF

Silicon Tracker



PCAL






FTOF

Drift Chambers
R1, R2, R3

Upgraded apparatus

	Forward detector	Central detector
Angular range Tracks Photons	5 – 40° 2.5 – 40°	35 – 125° n.a.
Resolution $\delta p/p$ $\delta\theta$ $\delta\phi$	< 1% @ 5 GeV/c < 1 mr < 3 mr	5% @ 1.5 GeV/c < 10-20 mr < 5 mr
Photon detection Energy $\delta\theta$	> 0.15 GeV 4 mr @ 1 GeV	n.a. n.a.
Neutron detection Efficiency	< 0.7	under dev.
Particle ID e/π π/p π/K K/p $\pi \rightarrow \gamma\gamma$ $\eta \rightarrow \gamma\gamma$	Full range Full range Full range < 4 GeV/c Full range Full range	n.a. < 1.25 GeV/c < 0.65 GeV/c < 1 GeV/c n.a. n.a.

Upgraded apparatus



GPD program

Proton DVCS

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS


Other highlights

GPD extraction

Summary

Proton DVCS A_{LU}


80 days @ $\mathcal{L} = 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 85% polarized beam



Statistical uncertainties from 1 % (low Q^2) to 10 % (high Q^2)


Proton DVCS A_{LU}

80 days @ $\mathcal{L} = 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 85% polarized beam



Statistical uncertainties from 1 % (low Q^2) to 10 % (high Q^2)

80 days @ $\mathcal{L} = 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 85% polarized beam



Dotted curve : no D-term, dashed-dotted : factorized t -dependence
 $Q^2 = 3.3 \text{ GeV}^2$, $x_B = 0.2$ (left and middle), $-t = 0.45 \text{ GeV}^2$ (left and right)

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

Proton DVCS A_{LU}

Generalized Parton
Distributions with
CLAS12

F.-X. Girod



Overview

Introduction

CLAS 12

Proposed
measurements


Proton DVCS

Neutron DVCS



Other highlights

GPD extraction

Summary



Extracted $H(\xi, \xi, t)$ and corresponding transverse profile



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS


Other highlights

GPD extraction

Summary

Proton DVCS TSA A_{UL}

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3



Statistical uncertainties from 2 % (low Q^2) to 10 % (high Q^2)

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS


Neutron DVCS

Other highlights

GPD extraction

Summary

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS


Neutron DVCS

Other highlights

GPD extraction

Summary

120 days @ $\mathcal{L} = 2 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with 80% polarized NH_3



Red solid line : $E = \tilde{E} = 0$, blue dashed line : $\tilde{H} = 0$

$Q^2 = 4.1 \text{ GeV}^2$, $x_B = 0.36$ (left and middle), $-t = 0.52 \text{ GeV}^2$ (left and right)

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

GPD program

Neutron DVCS

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS



Other highlights

GPD extraction

Summary

Neutron DVCS setup

For the detection of the scattered electron
and of the DVCS photon: CLAS12 +
Forward Calorimeter



For the detection of the recoil neutron:
Central Neutron Detector (CND)

Detection efficiency : 7 to 10 %

Acceptance for

charged particles:

- Central (CD), $40^\circ < \theta < 135^\circ$
- Forward (FD), $5^\circ < \theta < 40^\circ$

Acceptance for photons:

- FC $2.5^\circ < \theta < 5^\circ$
- EC, $5^\circ < \theta < 40^\circ$

Overview

Introduction

CLAS 12

Proposed
measurements


Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary



GPd program

Other highlights

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPd extraction

Summary

Transverse target asymmetries A_{UT} , DVCS & DVMP

More on angular momentum

Generalized Parton
Distributions with
CLAS12

F.-X. Girod



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS


Other highlights

GPD extraction

Summary

DVCS with frozen HD-ice


$Q^2=2.6, x_B=0.25$



$x_B \approx 0.25$

$Q^2 \approx 2.6 \text{ GeV}^2$

Exclusive ρ^0 production (hydrogen target)



$A_{UT} \sim \Delta_{\perp} \text{Im}AB^*$

$A \sim 2H^u + H^d$

$B \sim 2E^u + E^d$

GPD Extraction

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

Efforts towards GPD extraction



Overview

Introduction

CLAS 12

Proposed
measurements


Proton DVCS

Neutron DVCS


Other highlights

GPD extraction

Summary




H. Moutarde, **PRD 79** (2009) 094021



M. Guidal **PLB 689** (2010) 156

K. Kumericki & D. Mueller, arXiv:1008.2762



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights



GPD extraction

Summary

Conclusion

Summary and Prospects

- Deeply Virtual Compton Scattering on proton and neutron
- Flavor separation with Deeply Virtual Meson Production
- Valence quark tomography with global GPD extraction
- Also tomography of other hadrons
- An option of (polarized) positrons beam is also considered
- Sea and gluon tomography with Electron Ion Collider



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary


Additional slides

Mechanics and integration

Generalized Parton
Distributions with
CLAS12

F.-X. Girod

 Experimental Hall B



Overview

Introduction

CLAS 12

Proposed
measurements

Proton DVCS

Neutron DVCS

Other highlights

GPD extraction

Summary

Positron beam option

